

Lattice QCD at finite isospin chemical potential and temperature.

J. B. Kogut^a and D. K. Sinclair^{b *}

^aPhysics Department, University of Illinois, 1110 West Green Street, Urbana, IL 61801, USA

^bHEP Division, Argonne National Laboratory, 9700 South Cass Avenue, Argonne, IL 60439, USA

We simulate lattice QCD at a finite chemical potential μ_I for isospin (I_3) at zero and finite temperatures. At some $\mu_I = \mu_c$ QCD has a second order transition with mean-field critical exponents to a state where (I_3) is broken spontaneously by a charged pion condensate. Heating the system with $\mu_I > \mu_c$ we find there is some temperature at which this condensate evaporates. This transition appears to be second order and mean-field at lower μ_I values, and first order for μ_I sufficiently large. We are determining the dependence of the finite temperature crossover T_c on μ_I for $\mu_I < \mu_c$. This is expected to be identical to T_c 's dependence on quark-number chemical potential μ_q for small μ_q .

1. Introduction

QCD at finite baryon-number density describes nuclear matter including neutron stars and heavy nuclei. Nuclear matter at finite temperature may be observed in relativistic heavy-ion collisions at CERN and RHIC.

Because of the sign/phase difficulties associated with simulating QCD at finite baryon-number chemical potential, we have been studying related theories with positive fermion determinants. In particular, we are studying QCD (2-flavours) at finite chemical potential μ_I for isospin (I_3), at zero and finite temperature T . This represents a surface in the phase diagram for nuclear matter, which exists at finite isospin (I_3) density as well as finite baryon-number density.

From effective field theory (chiral perturbation theory) and other analyses, we expect a $T = 0$ transition at $\mu_I = m_\pi$ to a state with a charged pion condensate which breaks I_3 spontaneously, with the orthogonal charged pion state becoming a Goldstone boson [1] The transition is expected to be second order with mean-field critical exponents [1,2], a result which our simulations confirm [3].

For $\mu_I > m_\pi$ there is a finite T phase transition

where the pion condensate evaporates. For μ_I large enough, this transition appears to be first order. At low T the transition at $\mu_I = \mu_c \geq m_\pi$ appears to be second order and mean field as it is for $T = 0$ [3].

At $\mu_I = 0$ ($m \neq 0$) we know that there is a rapid crossover from ‘hadronic matter’ to a ‘quark-gluon’ plasma at $T = T_c \sim 150 - 200 \text{ MeV}$. This is the start of a line of such crossovers, which extends to finite μ_I . For small μ_I , the Swansea-Bielefeld collaboration have noted that $\beta_c(\mu_q) = \beta_c(\mu_I = 2\mu_q)$ to the extent that β_c is well defined for a crossover [4].

2. QCD at finite μ_I

The lattice quark action is

$$S_f = \sum_{\text{sites}} \left[\bar{\chi} \left[\not{D} \left(\frac{1}{2} \tau_3 \mu_I \right) + m \right] \chi + i \lambda \epsilon \bar{\chi} \tau_2 \chi \right] \quad (1)$$

which has a positive fermion determinant. λ is the explicit I_3 breaking parameter, required to observe spontaneous symmetry breaking on a finite lattice. We are interested in $\lambda \rightarrow 0$.

Our zero temperature simulations were performed on an 8^4 lattice at $m = 0.025$ and $m = 0.05$ at $\lambda = 0.1 \times m$ and $\lambda = 0.2 \times m$, using the hybrid molecular-dynamics method with ‘noisy’ fermions, and tuning to $N_f = 2$.

Figure 1 shows the charged pion condensate for the $m = 0.025$ runs, which shows evidence for a

*Talk presented by D. K. Sinclair. DKS was supported by DOE contract W-31-109-ENG-38. JBK was supported in part by an NSF grant NSF PHY-0102409. Simulations used IBM SPs and CRAYs at NERSC and NPACI.




Figure 1. Charged pion condensate as a function of I_3 chemical potential μ_I . The curves are a fit to mean-field scaling.

phase transition at $\mu_I = \mu_c$ to a state with a charged pion condensate at $\lambda = 0$. The fit is to a mean-field scaling form suggested by effective Lagrangian analyses.

For more details and measurements of other observables we refer the reader to [3].

3. QCD at finite μ_I and T

We have performed simulations on $8^3 \times 4$ lattices to study the finite temperature (T) behaviour of lattice QCD at finite μ_I [3].

Simulations at low β (4.0 and 5.0) and hence T have been performed, varying μ_I . Again we find that there is a phase transition for $\lambda = 0$ at some $\mu_c(\beta)$, above which the system is in a phase characterized by a charged pion condensate which breaks I_3 spontaneously. The transition appears to be second order with mean-field critical exponents, as in the $T = 0$ case.

We have also performed simulations at fixed $\mu_I > \mu_c(T = 0)$, increasing β (and hence T) until

the pion condensate ‘evaporates’ and the I_3 symmetry is restored. For large μ_I this phase transition is sufficiently abrupt to suggest that the transition has become first order. We are currently repeating these simulations on a larger ($16^3 \times 4$) lattice to check that the transition is indeed first order.




Figure 2. Pion condensate as a function of β for $m = 0.05$, $\lambda = 0.005$, and $\mu_I = 0.8$ on an $8^3 \times 4$ lattice.

Figure 2 shows the pion condensate for $\mu_I = 0.8$ as a function of β . The abrupt ‘jump’ between $\beta = 5.2$ and $\beta = 5.3$ suggests a first order transition. Preliminary simulations on a $16^3 \times 4$ lattice indicate that there is such an abrupt transition between $\beta = 5.265$ and $\beta = 5.270$.

4. The finite T transition at small μ_I

The finite temperature crossover ($m \neq 0$) from hadronic matter to a quark-gluon plasma extends to finite μ_I . We are running on $8^3 \times 4$ lattices with $m = 0.05$ and $N_f = 2$. For $\mu_I < \mu_c(T = 0) = 0.569(2)$ we set $\lambda = 0$.

At small μ_q , the Swansea-Bielefeld collaboration [4] found that the phase θ of the fermion determinant was ‘well-behaved’ – $\langle \cos(\theta) \rangle > 0$ and goes smoothly to 1 as $\mu_q \rightarrow 0$. As a consequence they expect, and find numerically by their series expansion methods, that $\beta_c(\mu_q) = \beta_c(\mu_I = 2\mu_q)$ to the extent that β_c is well defined for a crossover. Thus for small μ_I our simulations predict the small μ_q behaviour of β_c .

We have performed simulations at $\mu_I = 0.0, 0.1, 0.2, 0.3, 0.4$, and are currently running at $\mu_I = 0.5$, varying β through the crossover regime, measuring the chiral condensate, Wilson line, plaquette, isospin density and their susceptibilities, from which we will obtain the position of the crossover (the maxima of the susceptibilities) as a function of μ_I .




Figure 3. Chiral condensates for $\mu_I = 0.0, 0.1, 0.2, 0.3, 0.4$, as functions of β .

Figure 3 shows the chiral condensate measurements from these runs. We note that the temperature of the transition decreases *slowly* with increasing μ_I in qualitative agreement with the

μ_q behaviour reported by others [5,4,6,7].

5. Conclusions

At zero and low temperatures, there is some finite μ_I , μ_c say, above which I_3 is spontaneously broken by a charged pion condensate. This phase transition appears to be second order with mean-field critical exponents. Simulations will be performed on a $12^3 \times 24$ lattice to measure the spectrum of Goldstone and pseudo-Goldstone bosons, and to study instanton effects.

For $\mu_I > \mu_c$ there is a finite temperature phase transition at which the pion condensate evaporates. For large μ_I this transition appears to be first order. Simulations on a $16^3 \times 4$ lattice are being performed to check this.

We are studying the crossover from hadronic matter to a quark-gluon plasma at small μ_I and have confirmed that β_c and hence T_c decrease very slowly with increasing μ_I . This observation will be made quantitative. This is made more interesting because at low μ_I , the μ_I and μ_q dependence of this transition are believed to be identical. Simulations are planned to extend these studies of the μ_I dependence of the finite temperature transition to intermediate μ_I values. Larger-lattice simulations are needed.

It would be interesting to know if such charged pion condensates form when the system is also at finite baryon number density.

REFERENCES

1. D. T. Son and M. A. Stephanov, Phys. Rev. Lett. 86, 592 (2001); Yad. Fiz. 64, 899 (2001), Phys. Atom. Nucl. 64, 834 (2001).
2. K. Splittorff, D. Toublan and J. J. M. Verbaarschot, Nucl. Phys. B620, 290 (2002).
3. J. B. Kogut and D. K. Sinclair, hep-lat/0202028 (2002).
4. C. R. Allton, et al., hep-lat/0204010 (2002).
5. Z. Fodor and S. D. Katz, Phys. Lett. B534, 87 (2002); JHEP 03, 014 (2002).
6. P. de Forcrand and O. Philipsen, eprint hep-lat/0205016 (2002).
7. Talks presented by O. Philipsen, S. Ejiri, Z. Fodor, P. Crompton at this conference.